arXiv:hep-ph/0206118v2 29 Aug 2002

Renormalization Group Equation of Quark-Lepton Mass Matrices in the SO(10) Model with Two Higgs Scalars

Takeshi FUKUYAMA and Tatsuru KIKUCHI

Department of Physics, Ritsumeikan University, Kusatsu, Shiga, 525-8577 Japan

(February 1, 2008)

Abstract

The renormalization group equations (RGEs) of the mass matrices of quarks and leptons in a SO(10) model with two Higgs scalars in the Yukawa coupling are studied. This model is the minimal model of SUSY and non-SUSY SO(10) GUT which can reproduce all the experimental data. Non-SUSY SO(10) GUT model has the intermediate energy phase, Pati-Salam phase, and passes through the symmetry breaking pattern, $SO(10) \rightarrow SU(2)_L \times SU(2)_R \times SU(4)_C \rightarrow SU(2)_L \times U(1)_Y \times SU(3)_C$. Though minimal, it has, after the Pati-Salam phase, four Higgs doublets in Yukawa interactions. We consider the RGE's of the Yukawa coupling constants of quarks and charged leptons and of the coupling constants of the dimension five operators of neutrinos corresponding to the above symmetry breaking pattern. The scalar quartic interactions are also incorporated.

PACS number(s): 12.15.Ff, 12.10.-g, 12.60.-i

Typeset using REVTEX

In these decades many informations on the quark lepton mass matrices have been accumulated. We are confronted with the very era when we should seriously consider a realistic model in the scheme of grand unified theories. On this ocasion, neutrino masses may be the window to the grand unified theories via heavy right-handed neutrinos. Along this sight we considered in [1] and [2] the SO(10) model where two Higgs scalars participate in the Yukawa coupling. Our SO(10) model with two Higgs scalar, **10** and $\overline{\mathbf{126}}$, may be the first realistic model which successfully fit with the quark lepton mass spectra, Cabbibo-Kobayashi-Maskawa (CKM) [3], and Maki-Nakagawa-Sakata-Pontecorvo (MNSP) [4] mixing matrices based on the SO(10) framework. In these studies SO(10) invariant model is, of course, valid at the GUT scale and the data in our hand are those at much lower energy scale, at electroweak scale. So we must transport either of them to the other energy scale using the renormalization group equations (RGEs). In [5] we bottomed up the electroweak scale data about six quark masses, three mixing angles and one CP-phase in the CKM matrix and three charged lepton masses to the GUT scale and, using the SO(10) GUT mass relations (see Eqs.(7) and (18)) , obtained the neutrino mass matrix via seesaw mechanics [6] at GUT or at the intermediate scale. Then we toped down this mass matrix to the electroweak scale and checked whether it is consistent with the neutrino oscillation data and neutrinoless double beta decay. The results were very satisfactory in principle. However the theory is, of course, not conclusive. We adopted there some assumtions in order to extract the essential characters. One of them was the doublet-doblet splitting: that is, one of doublets is heavy relative to the other and our model in [5] was the same in essence as the MSSM except for the additional mixing angle among heavy and light Higgs doublets. However, this doublet-doublet splitting is not sure to occur. For instance, in the same SO(10) GUT with two Higgs scalar, these two Higgs doublet may work in parallel to remedy the over-abundance of the leptogenesis [7]. So in this paper we discarded this assumption and also incorporate scalar self coupling. In pay of this generalization we abandon in this paper the detailed data fitting and restrict ourselves in showing RGE's of quark-leptons mass matrices on more general ground. The renormalization of the neutrino mass operator was discussed by Babu-

Leung-Pantaleone and Chankowski-Pluciennik [8] for some simple models. Unfortunately it includes some calculational errors [9], though it does not affect our work, because we assume nuetrino couples to the Higgs doublets in a similar way to the up quarks. Our model is the most simple realistic model in SUSY and non-SUSY SO(10) GUT. At the GUT scale we have the Yukawa interactions to be given by

$$\mathcal{L}_Y = \sum_{i,j} \left(Y^{(10)}_{ij} \Psi^{iT} B \gamma^{\mu} H_{\mu} \Psi^j + \frac{1}{(5!)^2} Y^{(126)}_{ij} \Psi^{iT} B \gamma^{\mu_1}\gamma^{\mu_2}\gamma^{\mu_3}\gamma^{\mu_4}\gamma^{\mu_5} \overline{\Delta}_{\mu_1\mu_2\mu_3\mu_4\mu_5} \Psi^j + h.c. \right), \quad (1)$$

where $\Psi^i\,(i = 1, 2, 3)$ is the **16**-dimentional matter multiplet of the i-th generation, H and $\overline{\Delta}$ are the Higgs multiplet of **10** and $\overline{\mathbf{126}}$ representations under SO(10), respectively. B denotes the charge conjugation for SO(10) spinors : $B = \gamma_1\gamma_3\gamma_5\gamma_7\gamma_9$.

The Higgs scalar's quartic interactions with only H and $\overline{\Delta}$ are given by

$$\begin{aligned} V\left(H, \overline{\Delta}\right) &= \lambda_1 \left(H^{\mu} H_{\mu}\right)\left(H^{\nu} H_{\nu}\right) + \frac{1}{5!}\lambda_2 \left(H^{\mu} H_{\mu}\right)\left(\overline{\Delta}^{\mu_1\mu_2\mu_3\mu_4\mu_5}\overline{\Delta}_{\mu_1\mu_2\mu_3\mu_4\mu_5}\right) \\ &+ \frac{1}{4!}\lambda_3 \left(H^{\mu_1}\overline{\Delta}_{\mu_1\mu_2\mu_3\mu_4\mu_5}\right)\left(H_{\nu_1}\overline{\Delta}^{\nu_1\mu_2\mu_3\mu_4\mu_5}\right) \\ &+ \frac{1}{(5!)^2}\lambda_4 \left(\overline{\Delta}^{\mu_1\mu_2\mu_3\mu_4\mu_5}\overline{\Delta}_{\mu_1\mu_2\mu_3\mu_4\mu_5}\right)\left(\overline{\Delta}^{\nu_1\nu_2\nu_3\nu_4\nu_5}\overline{\Delta}_{\nu_1\nu_2\nu_3\nu_4\nu_5}\right) + h.c. \end{aligned} \quad (2)$$

Hereafter we consider non-SUSY SO(10) GUT explicitly. Gauge coupling unification needs the intermediate energy scale, Λ_I [10]. Between the grand unification scale and the intermediate scale, the effective Yukawa interactions are given by

$$-\mathcal{L}_Y = \sum_{i,j} \left(Y^{(10)}_{F\,ij} F_L^{iT} \Phi F_R^j + Y^{(126)}_{F\,ij} F_L^{iT} \Sigma F_R^j + Y^{(126)}_{R\,ij} F_R^{iT} \overline{\Delta_R} F_R^j + h.c. \right), \quad (3)$$

where F_L and F_R denote $(\mathbf{2}, \mathbf{1}, \mathbf{4})$ and $(\mathbf{1}, \mathbf{2}, \overline{\mathbf{4}})$ in Ψ^i, under $G_{224} \equiv SU(2)_L \times SU(2)_R \times SU(4)_C$, respectively. And also, Φ, Σ and $\overline{\Delta_R}$ correspond to $(\mathbf{2}, \mathbf{2}, \mathbf{1})$ in H, $(\mathbf{2}, \mathbf{2}, \mathbf{15})$ and $(\mathbf{1}, \mathbf{3}, \overline{\mathbf{10}})$ in $\overline{\Delta}$, respectively. Here we have assumed that suitably chosen $U(1)_H$ charge forbids the Yukawa interactions like, $F_L^{iT}\widetilde{\Phi}F_R^j$ and $F_L^{iT}\widetilde{\Sigma}F_R^j$, where

$$\widetilde{\Phi} = \epsilon^T \Phi^* \epsilon, \quad \widetilde{\Sigma} = \epsilon^T \Sigma^* \epsilon, \quad (4)$$

with

$$\epsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} . \tag{5}$$

And also the Higgs scalar's quartic interactions are given by

$$\begin{aligned} V(\Phi, \Sigma) &= \lambda_1 \left[\mathrm{tr}\left(\Phi \Phi^\dagger \right) \right]^2 + 2\lambda_2\, \mathrm{tr}\left(\Phi\Phi^\dagger \right) \mathrm{tr}\left(\Sigma\Sigma^\dagger \right) \\ &+ \lambda_3 \left\{ \left[\mathrm{tr}\left(\Phi\Sigma^\dagger \right) \right]^2 + \left[\mathrm{tr}\left(\Sigma\Phi^\dagger \right) \right]^2 \right\} + 4\lambda_4 \left[\mathrm{tr}\left(\Sigma\Sigma^\dagger \right) \right]^2 . \end{aligned} \tag{6}$$

When the intermediate symmetry breaking occures, $\overline{\Delta_R}$ have the vacuum expectation value and

$$M_R = \left\langle \overline{\Delta_R} \right\rangle Y_R^{(126)} . \tag{7}$$

Below the intermediate scale, it includes four Higgs doublets : two doublets (ϕ_1 and ϕ_3) come from $(\mathbf{2}, \mathbf{2}, \mathbf{1})$ and (ϕ_2 and ϕ_4) from $(\mathbf{2}, \mathbf{2}, \mathbf{15})$. $\{\phi_1,\ \phi_2\}$ couple with up-type quarks and leptons, and $\{\phi_3,\ \phi_4\}$ do with down-type quarks and leptons [11]. In these multiple Higgs models we always encounter with the flavor changing neutral current (FCNC) problem as Glashaw and Weinberg remarked [12]. However, by choosing the specific mass matrix textures, FCNC can be suppressed even in the case of multiple Higgs case [13]. Thus the low energy effective Yukawa interactions and the Higgs scalar's quartic interactions have the following forms,

$$\begin{aligned} -\mathcal{L}_Y = \sum_{i,j} \Big(& Y_{u\,ij}^{(10)}\, \overline{q_L^i}\, \widetilde{\phi_1} u_R^j + Y_{u\,ij}^{(126)}\, \overline{q_L^i}\, \widetilde{\phi_2} u_R^j \\ & + Y_{d\,ij}^{(10)}\, \overline{q_L^i}\, \phi_3 d_R^j + Y_{d\,ij}^{(126)}\, \overline{q_L^i}\, \phi_4 d_R^j \\ & + Y_{\nu\,ij}^{(10)}\, \overline{\ell_L^i}\, \widetilde{\phi_1} N_R^j + Y_{\nu\,ij}^{(126)}\, \overline{\ell_L^i}\, \widetilde{\phi_2} N_R^j \\ & + Y_{e\,ij}^{(10)}\, \overline{\ell_L^i}\, \phi_3 e_R^j + Y_{e\,ij}^{(126)}\, \overline{\ell_L^i}\, \phi_4 e_R^j + h.c. \Big) \\ & + \frac{1}{2} \sum_{i,j} \left(M_{R\,ij} \overline{N_R^{ci}} N_R^j + h.c. \right) , \end{aligned} \tag{8}$$

$$V(\phi) = \frac{1}{4!} \sum_{a,b.c.d=1,2,3,4} \lambda_{abcd} \left(\phi_a^\dagger \phi_b \right) \left(\phi_c^\dagger \phi_d \right) . \tag{9}$$

Here q_L and ℓ_L are the left-handed quark and lepton doublets :

$$q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix} \ , \quad \ell_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \ . \tag{10}$$

Also, ϕ_a are the Higgs doublets :

$$\phi_a = \begin{pmatrix} \phi_a^+ \\ \phi_a^0 \end{pmatrix} \ , \quad \widetilde{\phi}_a = \begin{pmatrix} \phi_a^{0*} \\ -\phi_a^- \end{pmatrix} \ . \tag{11}$$

From the hermiticity of (9), the self coupling constants λ_{abcd} satisfy

$$\lambda_{abcd} = \lambda_{cdab} = \lambda^*_{badc}. \tag{12}$$

Eq.(3) forbids the Yukawa and the scalar quartic interactions in Eqs.(8) and (9) which are not invariant under the $\mathbf{Z_2}$ symmetry,

$$\begin{aligned} &\phi_1 \to \phi_1, \ \ \phi_2 \to \phi_2, \\ &\phi_3 \to -\phi_3, \ \ \phi_4 \to -\phi_4 \end{aligned} \tag{13}$$

and corresponding transformations in the fermion sector. When the electroweak symmetry breaking occures, ϕ_a^0 $(a = 1, 2, 3, 4)$ have the vacuum expectation values(VEVs),

$$\left\langle \phi_a^0 \right\rangle = \frac{v_a}{\sqrt{2}} \quad (a = 1, 2, 3, 4) \tag{14}$$

and

$$\begin{aligned} M_u &= \frac{v_1^*}{\sqrt{2}} Y_u^{(10)} + \frac{v_2^*}{\sqrt{2}} Y_u^{(126)}, \\ M_d &= \frac{v_3}{\sqrt{2}} Y_d^{(10)} + \frac{v_4}{\sqrt{2}} Y_d^{(126)}, \\ M_D &= \frac{v_1^*}{\sqrt{2}} Y_\nu^{(10)} + \frac{v_2^*}{\sqrt{2}} Y_\nu^{(126)}, \\ M_e &= \frac{v_3}{\sqrt{2}} Y_e^{(10)} + \frac{v_4}{\sqrt{2}} Y_e^{(126)}. \end{aligned} \tag{15}$$

The Yukawa couplings above the intermediate energy scale are unified to

$$\begin{aligned} &\frac{1}{\sqrt{2}} Y_F^{(10)} \equiv Y^{(10)}, \\ &\frac{1}{4\sqrt{2}} Y_F^{(126)} = \frac{1}{4} Y_R^{(126)} \equiv Y^{(126)} \end{aligned} \tag{16}$$

at the GUT scale. Here the above numerical factors in Eq.(16) are necessarily to translate the $SO(10)$ language into the G_{224} language, see for example [14]. And also, at the the intermediate scale, the Yukawa couplings above the electroweak scale are unified to

$$\begin{aligned} &Y_u^{(10)} = Y_d^{(10)} = Y_\nu^{(10)} = Y_e^{(10)} \equiv Y_F^{(10)}, \\ &Y_u^{(126)} = Y_d^{(126)} = -\frac{1}{3} Y_\nu^{(126)} = -\frac{1}{3} Y_e^{(126)} \equiv Y_F^{(126)}, \end{aligned} \tag{17}$$

and Eq.(15) is reduced to [15]

$$\begin{aligned} M_u &= c_{10} M^{(10)} + c_{126} M^{(126)}, \quad M_d = M^{(10)} + M^{(126)}, \\ M_D &= c_{10} M^{(10)} - 3 c_{126} M^{(126)}, \quad M_e = M^{(10)} - 3 M^{(126)}. \end{aligned} \tag{18}$$

Here

$$c_{10} \equiv v_1^* / v_3 \,, \qquad c_{126} \equiv v_2^* / v_4 \,, \tag{19}$$

and

$$M^{(10)} \equiv \frac{v_3}{\sqrt{2}} Y_F^{(10)}, \quad M^{(126)} \equiv \frac{v_4}{\sqrt{2}} Y_F^{(126)}. \tag{20}$$

It goes from Eq.(6) that the Higgs quartic coupling constants are unified to

$$\begin{aligned} &\frac{\lambda_{abcd}}{4!} = \lambda_1 \ \text{ for } a,b,c,d = 1,3 \text{ only.} \\ &\frac{\lambda_{abcd}}{4!} = 2\lambda_2 \ \text{ for } a,b = 1,3 \text{ and } c,d = 2,4. \\ &\frac{\lambda_{abcd}}{4!} = \lambda_3 \ \text{ for } \{a,b\} \text{ and } \{c,d\} \text{ take any sets of } \{1,2\},\ \{1,4\},\ \{3,2\},\ \{3,4\}. \\ &\frac{\lambda_{abcd}}{4!} = 4\lambda_4 \ \text{ for } a,b,c,d = 2,4 \text{ only} \end{aligned} \tag{21}$$

at the intermediate scale. Here the coupling constants which does not satisfy Eqs.(2) are vanished and excluded from the above rule. It should be emphasized that this model can be compatible with the large angle of atmospheric neutrino oscillation as far as we do not adopt any simplification not allowed in SO(10) framework [2]. For $\Lambda_I \gg \mu$, M_R decouples and we must treat the four point interaction,

$$\begin{aligned}\mathcal{L}_{\nu\nu} &= \frac{1}{4}\sum_{a,b=1,2}\sum_{i,j}\kappa_{ij}^{(a,b)}\left(\overline{\ell_L^i}\widetilde{\phi_a}\right)\left(\phi^*_{b\alpha}\epsilon^{\alpha\beta}\ell^{Cj}_{L\beta}\right) + h.c. \\ &= -\frac{1}{2}\sum_{a,b=1,2}\sum_{i,j}\kappa_{ij}^{(a,b)}\left\{\left(\overline{\nu_L^i}\nu_L^{Cj}\phi_a^{0*}\phi_b^{0*} + \overline{e_L^i}e_L^{Cj}\phi_a^-\phi_b^-\right)\right. \\ &\quad\left. - \frac{1}{2}\left(\overline{\nu_L^i}e_L^{Cj} + \overline{e_L^i}\nu_L^{Cj}\right)\left(\phi_a^-\phi_b^{0*} + \phi_a^{0*}\phi_b^-\right)\right\} + h.c., \end{aligned} \tag{22}$$

where i, j are flavour indicies. The reason why a, b run over $1, 2$ is that it comes from the Dirac neutrinos (the third line in the right-hand side of Eq.(5)). Thus the effective light neutrino mass matrix is given by

$$M_\nu = \frac{1}{2}\sum_{a,b=1,2}\kappa^{(a,b)}v_a^* v_b^*. \tag{23}$$

We have two Higgs doublets in each up quarks, down quarks, Dirac neutrinos and charged leptons and $\kappa^{(a,b)}$ $(a, b = 1, 2)$ in ours are the generaslization of $\kappa^{(22)}$ in [8]. We have no term corresponding to $\kappa^{(11)}$ in [8].

In the non SUSY SO(10) with two Higgs scalars, the one loop RGEs for the effective Yukawa couplings first at the energy region between the grand unification scale and the intermediate sale are given by :

$$\begin{aligned}16\pi^2\frac{dY_F^{(10)}}{dt} &= \left(Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}Y_F^{(126)}Y_F^{(126)\dagger}\right)Y_F^{(10)} \\ &\quad + Y_F^{(10)}\left\{Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}\left(Y_F^{(126)}Y_F^{(126)\dagger} + Y_R^{(126)}Y_R^{(126)\dagger}\right)\right\} \\ &\quad + 4\mathrm{tr}\left(Y_F^{(10)}Y_F^{(10)\dagger}\right)Y_F^{(10)} + \left(\frac{9}{4}g_{2L}^2 + \frac{9}{4}g_{2R}^2 + \frac{15}{4}g_{4C}^2\right)Y_F^{(10)}, \end{aligned} \tag{24}$$

$$\begin{aligned}16\pi^2\frac{dY_F^{(126)}}{dt} &= \left(Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}Y_F^{(126)}Y_F^{(126)\dagger}\right)Y_F^{(126)} \\ &\quad + Y_F^{(126)}\left\{Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}\left(Y_F^{(126)}Y_F^{(126)\dagger} + Y_R^{(126)}Y_R^{(126)\dagger}\right)\right\} \\ &\quad + \mathrm{tr}\left(Y_F^{(126)}Y_F^{(126)\dagger}\right)Y_F^{(126)} + \left(\frac{9}{4}g_{2L}^2 + \frac{9}{4}g_{2R}^2 + \frac{15}{4}g_{4C}^2\right)Y_F^{(126)}, \end{aligned} \tag{25}$$

$$\begin{aligned}16\pi^2\frac{dY_R^{(126)}}{dt} &= \left\{Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}\left(Y_F^{(126)}Y_F^{(126)\dagger} + Y_R^{(126)}Y_R^{(126)\dagger}\right)\right\}Y_R^{(126)} \\ &\quad + Y_R^{(126)}\left\{Y_F^{(10)}Y_F^{(10)\dagger} + \frac{15}{4}\left(Y_F^{(126)}Y_F^{(126)\dagger} + Y_R^{(126)}Y_R^{(126)\dagger}\right)\right\} \\ &\quad + \mathrm{tr}\left(Y_R^{(126)}Y_R^{(126)\dagger}\right)Y_R^{(126)} + \left(\frac{9}{2}g_{2R}^2 + \frac{15}{4}g_{4C}^2\right)Y_R^{(126)}, \end{aligned} \tag{26}$$

where g_{2L}, g_{2R} and g_{4C} are the $SU(2)_L$, $SU(2)_R$ and $SU(4)_C$ gauge coupling constants, respectively. At the second stage, the energy region between the intermediate sale and the weak scale, the one loop RGEs for the effective Yukawa couplings are given by :

$$
\begin{aligned}
16\pi^2 \frac{dY_u^{(10)}}{dt} &= 3\mathrm{tr}(Y_u^{(10)} Y_u^{(10)\dagger}) Y_u^{(10)} + 3\mathrm{tr}(Y_u^{(10)} Y_u^{(126)\dagger}) Y_u^{(126)} \\
&- \left(8g_3^2 + \frac{9}{4}g_2^2 + \frac{17}{12}g_Y^2\right) Y_u^{(10)} \\
&+ \frac{1}{2}\left(Y_u^{(10)} Y_u^{(10)\dagger} + Y_u^{(126)} Y_u^{(126)\dagger}\right. \\
&+ \left. Y_d^{(10)} Y_d^{(10)\dagger} + Y_d^{(126)} Y_d^{(126)\dagger}\right) Y_u^{(10)} \\
&+ Y_u^{(10)}\left(Y_u^{(10)\dagger} Y_u^{(10)} + Y_u^{(126)\dagger} Y_u^{(126)}\right), \qquad (27) \\
16\pi^2 \frac{dY_u^{(126)}}{dt} &= 3\mathrm{tr}(Y_u^{(126)} Y_u^{(126)\dagger}) Y_u^{(126)} + 3\mathrm{tr}(Y_u^{(126)} Y_u^{(10)\dagger}) Y_u^{(10)} \\
&- \left(8g_3^2 + \frac{9}{4}g_2^2 + \frac{17}{12}g_Y^2\right) Y_u^{(126)} \\
&+ \frac{1}{2}\left(Y_u^{(10)} Y_u^{(10)\dagger} + Y_u^{(126)} Y_u^{(126)\dagger}\right. \\
&+ \left. Y_d^{(10)} Y_d^{(10)\dagger} + Y_d^{(126)} Y_d^{(126)\dagger}\right) Y_u^{(126)} \\
&+ Y_u^{(126)}\left(Y_u^{(10)\dagger} Y_u^{(10)} + Y_u^{(126)\dagger} Y_u^{(126)}\right), \qquad (28) \\
16\pi^2 \frac{dY_d^{(10)}}{dt} &= \left\{3\mathrm{tr}(Y_d^{(10)} Y_d^{(10)\dagger}) + \mathrm{tr}(Y_e^{(10)} Y_e^{(10)\dagger})\right\} Y_d^{(10)} \\
&+ \left\{3\mathrm{tr}(Y_d^{(10)} Y_d^{(126)\dagger}) + \mathrm{tr}(Y_e^{(10)} Y_e^{(126)\dagger})\right\} Y_d^{(126)} \\
&- \left(8g_3^2 + \frac{9}{4}g_2^2 + \frac{5}{12}g_Y^2\right) Y_d^{(10)} \\
&+ \frac{1}{2}\left(Y_u^{(10)} Y_u^{(10)\dagger} + Y_u^{(126)} Y_u^{(126)\dagger}\right. \\
&+ \left. Y_d^{(10)} Y_d^{(10)\dagger} + Y_d^{(126)} Y_d^{(126)\dagger}\right) Y_d^{(10)} \\
&+ Y_d^{(10)}\left(Y_d^{(10)\dagger} Y_d^{(10)} + Y_d^{(126)\dagger} Y_d^{(126)}\right), \qquad (29) \\
16\pi^2 \frac{dY_d^{(126)}}{dt} &= \left\{3\mathrm{tr}(Y_d^{(126)} Y_d^{(126)\dagger}) + \mathrm{tr}(Y_e^{(126)} Y_e^{(126)\dagger})\right\} Y_d^{(126)} \\
&+ \left\{3\mathrm{tr}(Y_d^{(126)} Y_d^{(10)\dagger}) + \mathrm{tr}(Y_e^{(126)} Y_e^{(10)\dagger})\right\} Y_d^{(10)} \\
&- \left(8g_3^2 + \frac{9}{4}g_2^2 + \frac{5}{12}g_Y^2\right) Y_d^{(126)} \\
&+ \frac{1}{2}\left(Y_u^{(10)} Y_u^{(10)\dagger} + Y_u^{(126)} Y_u^{(126)\dagger}\right. \\
&+ \left. Y_d^{(10)} Y_d^{(10)\dagger} + Y_d^{(126)} Y_d^{(126)\dagger}\right) Y_d^{(126)}
\end{aligned}
$$

$$
\begin{aligned}
&+ Y_d^{(126)}\left(Y_d^{(10)\dagger}Y_d^{(10)} + Y_d^{(126)\dagger}Y_d^{(126)}\right), \qquad (30)\\
16\pi^2\frac{dY_e^{(10)}}{dt} &= \left\{3\mathrm{tr}(Y_d^{(10)}Y_d^{(10)\dagger}) + \mathrm{tr}(Y_e^{(10)}Y_e^{(10)\dagger})\right\}Y_e^{(10)}\\
&+ \left\{3\mathrm{tr}(Y_d^{(10)}Y_d^{(126)\dagger}) + \mathrm{tr}(Y_e^{(10)}Y_e^{(126)\dagger})\right\}Y_e^{(126)}\\
&- \left(\frac{9}{4}g_2^2 + \frac{15}{4}g_Y^2\right)Y_e^{(10)}\\
&+ \frac{1}{2}\left(Y_e^{(10)}Y_e^{(10)\dagger} + Y_e^{(126)}Y_e^{(126)\dagger}\right)Y_e^{(10)}\\
&+ Y_e^{(10)}\left(Y_e^{(10)\dagger}Y_e^{(10)} + Y_e^{(126)\dagger}Y_e^{(126)}\right), \qquad (31)\\
16\pi^2\frac{dY_e^{(126)}}{dt} &= \left\{3\mathrm{tr}(Y_d^{(126)}Y_d^{(126)\dagger}) + \mathrm{tr}(Y_e^{(126)}Y_e^{(126)\dagger})\right\}Y_e^{(126)}\\
&+ \left\{3\mathrm{tr}(Y_d^{(126)}Y_d^{(10)\dagger}) + \mathrm{tr}(Y_e^{(126)}Y_e^{(10)\dagger})\right\}Y_e^{(10)}\\
&- \left(\frac{9}{4}g_2^2 + \frac{15}{4}g_Y^2\right)Y_e^{(126)}\\
&+ \frac{1}{2}\left(Y_e^{(10)}Y_e^{(10)\dagger} + Y_e^{(126)}Y_e^{(126)\dagger}\right)Y_e^{(126)}\\
&+ Y_e^{(126)}\left(Y_e^{(10)\dagger}Y_e^{(10)} + Y_e^{(126)\dagger}Y_e^{(126)}\right). \qquad (32)
\end{aligned}
$$

In Eqs.(27)-(32), the first two terms (before g_i^2 terms) are the contribution of fermion loop, the third term that of gauge loop, and the remaining two terms those of Higgs loop. The second term indicates the mixing of two Higgs doublets in the fermion loop correction. These formulas, of course, are reduced to one-loop RGEs for the standard model if we set either $Y^{(10)}$ or $Y^{(126)}$ zero [16]. The one loop RGEs for the scalar quartic couplings in our model are [17]

$$
\begin{aligned}
16\pi^2\frac{d\lambda_{abcd}}{dt} &= \frac{1}{6}\sum_{m,n=1,2,3,4}(2\lambda_{abmn}\lambda_{nmcd} + \lambda_{abmn}\lambda_{cmnd} + \lambda_{amnb}\lambda_{mncd}\\
&+ \lambda_{amnd}\lambda_{cnmb} + \lambda_{amcn}\lambda_{mbnd}) - 3\left(3g_2^2 + g_Y^2\right)\lambda_{abcd}\\
&+ 9\left(3g_2^4 + g_Y^4\right)\delta_{ab}\delta_{cd} + 36\,g_2^2g_Y^2\left(\delta_{ad}\delta_{bc} - \frac{1}{2}\delta_{ab}\delta_{cd}\right)\\
&+ \sum_{m,n=1,2,3,4}(\lambda_{mbcd}A_{am} + \lambda_{amcd}A_{mb} + \lambda_{abmd}A_{cm} + \lambda_{abcm}A_{md})\\
&- 48\,H_{abcd} \quad (a,b,c,d = 1,2,3,4), \qquad (33)
\end{aligned}
$$

Here

$$
A_{ab} \equiv \mathrm{tr}\left(3Y_a^{u\dagger}Y_b^u + 3Y_a^{d\dagger}Y_b^d + Y_a^{e\dagger}Y_b^e\right), \qquad (34)
$$

and

$$H_{abcd} \equiv \mathrm{tr}\left(3Y_d^{u\dagger}Y_c^uY_b^{u\dagger}Y_a^u + 3Y_a^{d\dagger}Y_b^dY_c^{d\dagger}Y_d^d + Y_a^{e\dagger}Y_b^eY_c^{e\dagger}Y_d^e \right.$$
$$\left. + 3Y_a^{u\dagger}Y_b^uY_d^{d\dagger}Y_c^d + 3Y_b^{d\dagger}Y_a^dY_c^{u\dagger}Y_d^u - 3Y_d^{d\dagger}Y_c^dY_b^{u\dagger}Y_a^u - 3Y_a^{u\dagger}Y_d^uY_b^{d\dagger}Y_c^d\right), \tag{35}$$

with

$$Y_1^u = Y_u^{(10)}, \quad Y_2^u = Y_u^{(126)}, \quad Y_3^d = Y_d^{(10)}, \quad Y_4^d = Y_d^{(126)},$$
$$Y_3^e = Y_e^{(10)}, \quad Y_4^e = Y_e^{(126)}, \quad \text{and otherwise zero.} \tag{36}$$

As for the RGEs of VEV's of Higgs fields, there may be some conflicts. Someone consider them constants [5] [18] [19], and other ones make them evolve as $\frac{d\sqrt{-2\mu^2/\lambda}}{dt}$ for the simplest case [20]. The situation may depend on what scale and what object we consider. Here we adopt the standpoint that the RGEs of v_a are those of ϕ_a^0 [16] [21] [22]. That is, in our case,

$$16\pi^2\frac{dv_1}{dt} = -3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(10)\dagger}\right)v_1 - 3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(126)\dagger}\right)v_2$$
$$+ \left(\frac{9}{4}g_2^2 + \frac{3}{4}g_Y^2\right)v_1, \tag{37}$$

$$16\pi^2\frac{dv_2}{dt} = -3\mathrm{tr}\left(Y_u^{(126)}Y_u^{(126)\dagger}\right)v_2 - 3\mathrm{tr}\left(Y_u^{(126)}Y_u^{(10)\dagger}\right)v_1$$
$$+ \left(\frac{9}{4}g_2^2 + \frac{3}{4}g_Y^2\right)v_2, \tag{38}$$

$$16\pi^2\frac{dv_3}{dt} = -\mathrm{tr}\left(3Y_d^{(10)\dagger}Y_d^{(10)} + Y_e^{(10)\dagger}Y_e^{(10)}\right)v_3$$
$$- \mathrm{tr}\left(3Y_d^{(10)\dagger}Y_d^{(126)} + Y_e^{(10)\dagger}Y_e^{(126)}\right)v_4$$
$$+ \left(\frac{9}{4}g_2^2 + \frac{3}{4}g_Y^2\right)v_3, \tag{39}$$

$$16\pi^2\frac{dv_4}{dt} = -\mathrm{tr}\left(3Y_d^{(126)\dagger}Y_d^{(126)} + Y_e^{(126)\dagger}Y_e^{(126)}\right)v_4$$
$$- \mathrm{tr}\left(3Y_d^{(126)\dagger}Y_d^{(10)} + Y_e^{(126)\dagger}Y_e^{(10)}\right)v_3$$
$$+ \left(\frac{9}{4}g_2^2 + \frac{3}{4}g_Y^2\right)v_4. \tag{40}$$

Here we have replaced the RGEs of ϕ_a with v_a. It goes from Eq.(15) and Eqs.(27)-(40) that

$$16\pi^2\frac{dM_u}{dt} = -\frac{3}{\sqrt{2}}\,\mathrm{tr}\left(Y_u^{(10)}Y_u^{(126)\dagger} - Y_u^{(126)}Y_u^{(10)\dagger}\right)\left(v_2Y_u^{(10)} - v_1Y_u^{(126)}\right)$$
$$- \left(8g_3^2 + \frac{2}{3}g_Y^2\right)M_u$$

$$
\begin{aligned}
&+\frac{1}{2}\left(Y_u^{(10)}Y_u^{(10)\dagger}+Y_u^{(126)}Y_u^{(126)\dagger}\right.\\
&+\left.Y_d^{(10)}Y_d^{(10)\dagger}+Y_d^{(126)}Y_d^{(126)\dagger}\right)M_u\\
&+M_u\left(Y_u^{(10)\dagger}Y_u^{(10)}+Y_u^{(126)\dagger}Y_u^{(126)}\right),
\end{aligned}
\tag{41}
$$

$$
\begin{aligned}
16\pi^2\frac{dM_d}{dt}&=-\left(8g_3^2-\frac{1}{3}g_Y^2\right)M_d\\
&+\frac{1}{2}\left(Y_u^{(10)}Y_u^{(10)\dagger}+Y_u^{(126)}Y_u^{(126)\dagger}\right.\\
&+\left.Y_d^{(10)}Y_d^{(10)\dagger}+Y_d^{(126)}Y_d^{(126)\dagger}\right)M_d\\
&+M_d\left(Y_d^{(10)\dagger}Y_d^{10}+Y_d^{(126)\dagger}Y_d^{126}\right),
\end{aligned}
\tag{42}
$$

$$
\begin{aligned}
16\pi^2\frac{dM_e}{dt}&=-3g_Y^2M_e\\
&+\frac{1}{2}\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)M_e\\
&+M_e\left(Y_e^{(10)\dagger}Y_e^{(10)}+Y_e^{(126)\dagger}Y_e^{(126)}\right).
\end{aligned}
\tag{43}
$$

It should be remarked that RGEs destroy the transpose-invariance of mass matrix possesed at GUT. At the region $\Lambda_I \gg \mu$, we must treat the RGEs of κ of Eq.(22), giving the following forms;

$$
\begin{aligned}
16\pi^2\frac{d\kappa^{(1,1)}}{dt}&=6\mathrm{tr}\left(Y_u^{(10)}Y_u^{(10)\dagger}\right)\kappa^{(1,1)}\\
&+3\mathrm{tr}\left(Y_u^{(126)}Y_u^{(10)\dagger}\right)\left(\kappa^{(1.2)}+\kappa^{(2.1)}\right)\\
&-3g_2^2\kappa^{(1,1)}\\
&+\frac{1}{6}\left(\lambda_{1111}\kappa^{(1,1)}+\lambda_{1112}\kappa^{(1,2)}+\lambda_{1212}\kappa^{(2,2)}\right)\\
&+\frac{1}{2}\left\{\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)\kappa^{(1,1)}\right.\\
&+\left.\kappa^{(1,1)}\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)^T\right\},
\end{aligned}
\tag{44}
$$

$$
\begin{aligned}
16\pi^2\frac{d\kappa^{(2,2)}}{dt}&=6\mathrm{tr}\left(Y_u^{(126)}Y_u^{(126)\dagger}\right)\kappa^{(2,2)}\\
&+3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(126)\dagger}\right)\left(\kappa^{(1.2)}+\kappa^{(2,1)}\right)\\
&-3g_2^2\kappa^{(2,2)}\\
&+\frac{1}{6}\left(\lambda_{2222}\kappa^{(2,2)}+\lambda_{2121}\kappa^{(1,1)}\right)\\
&+\frac{1}{2}\left\{\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)\kappa^{(2,2)}\right.
\end{aligned}
$$

$$+\ \kappa^{(2,2)}\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)^T\Bigg\}, \tag{45}$$

$$\begin{aligned}
16\pi^2\frac{d\kappa^{(1,2)}}{dt} &= 3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(10)\dagger}+Y_u^{(126)}Y_u^{(126)\dagger}\right)\kappa^{(1,2)} \\
&+ 3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(126)\dagger}\right)\kappa^{(1.1)} \\
&+ 3\mathrm{tr}\left(Y_u^{(126)}Y_u^{(10)\dagger}\right)\kappa^{(2.2)} \\
&- g_2^2\left(2\kappa^{(1,2)}+\kappa^{(2,1)}\right) \\
&+ \frac{1}{6}\left(\lambda_{1122}\kappa^{(1,2)}+\lambda_{1221}\kappa^{(2,1)}\right) \\
&+ \frac{1}{2}\Bigg\{\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)\kappa^{(1,2)} \\
&+\ \kappa^{(1,2)}\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)^T\Bigg\}, \qquad (46) \\
16\pi^2\frac{d\kappa^{(2,1)}}{dt} &= 3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(10)\dagger}+Y_u^{(126)}Y_u^{(126)\dagger}\right)\kappa^{(2,1)} \\
&+ 3\mathrm{tr}\left(Y_u^{(10)}Y_u^{(126)\dagger}\right)\kappa^{(1.1)} \\
&+ 3\mathrm{tr}\left(Y_u^{(126)}Y_u^{(10)\dagger}\right)\kappa^{(2.2)} \\
&- g_2^2\left(\kappa^{(1,2)}+2\kappa^{(2,1)}\right) \\
&+ \frac{1}{6}\left(\lambda_{2211}\kappa^{(2,1)}+\lambda_{2112}\kappa^{(1,2)}\right) \\
&+ \frac{1}{2}\Bigg\{\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)\kappa^{(2,1)} \\
&+\ \kappa^{(2,1)}\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)^T\Bigg\}, \qquad (47)
\end{aligned}$$

The different factor 2 of the coefficients of g_2^2 in $\kappa^{(1,2)}$ and $\kappa^{(2,1)}$ comes from the different contribution of W-boson and Z-boson loop correction on $\kappa^{(1,2)}$ and $\kappa^{(2,1)}$. The self coupling contributions come from the diagram Fig.1.

Substituting the above equations into Eq.(18), we obtain the RGE of the light neutrino mass matrix:

$$\begin{aligned}
16\pi^2\frac{dM_\nu}{dt} &= \frac{1}{2}\Bigg\{3(g_2^2+g_Y^2)M_\nu \\
&+ \frac{1}{6}\sum_{a,b,c,d=1,2}\lambda_{abcd}v_a^*v_c^*\kappa^{(b,d)} \\
&+ \left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)M_\nu \\
&+ M_\nu\left(Y_e^{(10)}Y_e^{(10)\dagger}+Y_e^{(126)}Y_e^{(126)\dagger}\right)^T\Bigg\}. \qquad (48)
\end{aligned}$$

Thus we have obtained RGEs of SO(10) GUT with two Higgs scalars. These formulas are especially important for essentially multiple Higgs models where two Higgs doblets in up or down type quark-lepton mass matrices can not suffer doublet-doublet splittings. Even in a model where the doublet-doublet splitting occurs, our results are useful for estimating the threhold effects due to the heavy pairs of Higgs doublets. We calculated RGE's explicitly in non-SUSY SO(10). Those of SUSY SO(10) are obtained analogously by considering the contributions of the superpartners of non-SUSY contents except for the scalar self couplings.

ACKNOWLEDGMENTS

We thank N.Okada, Y.Koide, and K.Matsuda for very useful arguments. T.F expresses his gratitude to R.Mohapatra for hospitality at the University of Maryland where a large part of this work was accomplished.

REFERENCES

[1] K. Matsuda, Y. Koide, T. Fukuyama, Phys. Rev. **D64**, 053015 (2001).

[2] K. Matsuda, Y. Koide, T. Fukuyama and H. Nishiura, Phys. Rev. **D65**, 033008 (2002).

[3] N. Cabibbo, Phys.Rev.Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

[4] Z. Maki, M. Nakagawa, and S. Sakata, Prog.Theor.Phys. **28**, 247 (1962); B. Pontecorvo, J. Exptl. Theoret. Phys., **33**, 549 (1957) [Sov. Phys. JETP **6**, 429 (1958)]; J. Exptl. Theoret. Phys., **34**, 247 (1958) [Sov. Phys. JETP **7**, 172 (1958)].

[5] T. Fukuyama and N. Okada, UMD-PP-02-048, May 2002. 17pp. [Archive:hep-ph/0205066].

[6] T. Yanagida,in Proceedings of the workshop on the Unified Theory and Baryon Number in the Universe, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979); M. Gell-Mann, P. Ramond, and R. Slansky, in Supergravity, edited by D. Freedman and P.van Niewenhuizen (north-Holland, Amsterdam 1979);R.N. Mohapatra and G. Senjanovic, Phys.Rev.Lett. **44**, 912 (1980).

[7] T. Fukuyama and N. Okada, UMD-PP-02-036, Feb 2002. 11pp. [Archive:hep-ph/0202214].

[8] K.S. Babu, C.N. Leung, and J. Pantaleone, Phys. Lett. **B319**, 191 (1993); P.H. Chankowski and Z. Pluciennik, Phys. Lett. **B316**, 312 (1993).

[9] S. Antusch, M. Drees, J. Kersten, M. Lindner, and M. Ratz, Phys. Lett. **B519**, 238 (2001); Phys. Lett. **B525**, 130 (2002).

[10] J.C. Pati and A. Salam, Phys.Rev. **D10**, 275 (1974).

[11] K. Matsuda, T. Fukuyama and H. Nishiura, Phys.Rev. **D61**, 053001 (2000).

[12] S.L. Glashow and S. Weinberg, Phys. Rev. **D15**, 1958 (1977).

[13] T.P. Cheng, M. Sher, Phys.Rev. **D35**, 3484 (1987); A. Antaramian, L.J. Hall, A. Rasin, Phys.Rev.Lett. **69**, 1871-1873 (1992).

[14] C.S. Aulakh, A. Girdhar, [Archive: hep-ph/0204097].

[15] K.S. Babu and R.N. Mohapatra, Phy. Rev. Lett. **70**, 2845 (1993); D-G. Lee and R.N. Mohapatra, Phys. Rev. **D51**, 1353 (1995).

[16] H. Arason et al., Phys. Rev **D46**, 3945 (1992).

[17] T.P. Cheng, E. Eichten and Ling-Fong Li, Phys. Rev. **D9**, 2259 (1974); H. Komatsu, Prog. Theor. Phys. **67**, 1177, (1982); M.E. Machacek and M.T. Vaughn, Nucl. Phys. **B249**, 70 (1985).

[18] H. Fusaoka and Y. Koide, Phys. Rev. **D57**, 3986 (1998).

[19] J. Ellis and S. Lola, Phys. Lett. **B458**, 310 (1999).

[20] P.H. Chankowski and P. Wasowicz, Eur.Phys.J. **C23**, 249-258 (2002).

[21] G. Cvetic, S.S. Hwang and C.S. Kim, Phys. Rev. **D58**, 116003 (1998).

[22] C.R. Das and M.K. Parida, Eur. Phys. J **C20**, 121 (2001).

FIGURES



FIG. 1. The Higgs scalar self couplings contribution to the RGE's of κ.